



User name: _____
Password: _____
[Register](#) [Forgotten Password](#) [Athens Login](#)

[Synergy Home](#) | [Browse](#) | [Search](#) | [My Synergy](#) | [Books Online](#) | [Resources](#) | [About](#) | [Help](#)



JOURNAL OF Veterinary Pharmacology and Therapeutics

Journal Menu

[Table of Contents](#)
[List of Issues](#)

Tools

[Email this article](#)
[Add to favorite articles](#)
[Export this citation](#)
[Alert me when this article is cited: Email | RSS](#)
(What is this?)

[View PubMed citation](#)
[View ISI citation](#)

Publication history

Issue online:
05 Jan 2002

[Home](#) > [List of Issues](#) > [Table of Contents](#) > [Article Abstract](#)


Journal of Veterinary Pharmacology and Therapeutics

Volume 21 Issue 1 Page 74 - February 1998

To cite this article: ELLIOTT, BRADY (1998)

Characterization of vasodilatory adenosine receptors in equine digital veins

Journal of Veterinary Pharmacology and Therapeutics 21 (1), 74-81.
doi:10.1046/j.1365-2885.1998.00106.x



[Prev Article](#) | [Next Article](#)

Original Article

Characterization of vasodilatory adenosine receptors in equine digital veins

ELLIOTT & BRADY

¹Department of Veterinary Basic Sciences, Royal Veterinary College,
Royal College Street, London NW1 0TU, UK

Correspondence to: Elliott Department of Veterinary Basic
Sciences, Royal Veterinary College, Royal College Street,
London NW1 0TU, UK

Abstract

Isolated equine digital veins (EDVs) which had been denuded of their endothelium were used to study adenosine receptors causing vasodilation. When the blood vessel wall tension was raised with the thromboxanemimetic, U44069 (30 nM), the order of vasodilator potency of adenosine receptor agonists was: 5'-N-ethylcarboxamidoadenosine (NECA) > 2-*p*-(2-carboxyethyl)phenyl amino-5'-N-ethylcarboxamido-adenosine (CGS 21680) > 5'-N-methylcarboxamido-adenosine (MECA) >> N⁶-cyclohexyladenosine (CHA) > N⁶-cyclopentyladenosine (CPA) > N⁶-2-(4-Aminophenyl) ethyladenosine (APNEA) > adenosine. Removal of the endothelium had no significant effect on the responses to NECA. The adenosine receptor antagonists, 8-cyclopentyl-1,3-dipropylxanthine (DPCPX; A₁-selective) and xanthine amine cogener (XAC; non-selective

J Neurophysiol 79: 501-510, 1998;
0022-3077/98 \$5.00

The Journal of Neurophysiology Vol. 79 No. 2 February 1998, pp. 501-510
Copyright ©1998 by the American Physiological Society

Adenosine A1 Receptors Mediate Retinotectal Presynaptic Inhibition: Uncoupling by C-Kinase and Role in LTP During Regeneration

Chunyi Zhang and John T. Schmidt

Department of Biological Sciences and Neurobiology Research Center,
State University of New York, Albany, New York 12222

► ABSTRACT

Zhang, Chunyi and John T. Schmidt. Adenosine A1 receptors mediate retinotectal presynaptic inhibition: uncoupling by C-kinase and role in LTP during regeneration. *J. Neurophysiol.* 79: 501-510, 1998. Presynaptic adenosine receptors inhibit transmitter release at many synapses and are known to

exist on retinotectal terminals. In this paper we show that adenosine decreases retinotectal field potentials by ~30% and investigate the mechanism. First, as judged by the effects of specific calcium channel blockers, retinotectal transmission was mediated almost exclusively by N-type calcium channels, which are known to be modulated by adenosine A1 receptors. Transmission was completely blocked by either ω -Conotoxin GVIA (–100%, N-type blocker) or ω -Conotoxin MVIIC (–99%, N-, P- and Q-type blocker) and was not significantly affected by ω -Agatoxin IVA [$+1.7 \pm 9.3\%$ (SE), P-, Q-type blocker], but was augmented slightly by nifedipine ($+9.3 \pm 2.1\%$, L-type blocker). Second, the adenosine inhibition was presynaptic, as indicated by a 43% increase in paired-pulse facilitation. Third, the selective A1 agonist cyclohexyl adenosine (CHA) at 50 nM caused a 21% decrease in amplitude and the selective A2 agonist *N*⁶-[2-(3,5-dimethoxyphenyl)-2-(2-methylphenyl)-ethyl]adenosine (DPMA) at 100 nM caused a 24% increase. Fourth, the selective A1 antagonist 8-cyclopentyl-1,3-dipropylxanthine (DPCPX) alone produced an increase in the field potential, suggesting a tonic inhibition mediated by endogenous adenosine. Fifth, pertussis toxin eliminated adenosine inhibition implicating G_i or G_o protein coupling. Sixth, C-kinase activation eliminated the A1-mediated inhibition. In regenerating projections, adenosine also caused a decrease in transmission ($-30 \pm 12\%$), but after induction of long-term potentiation (LTP) via trains of stimuli or via treatment with the phosphatase inhibitor okadaic acid, the adenosine response was converted to an augmentation. Because LTP is associated with C-kinase activation, this is consistent with C-kinase uncoupling the A1 receptor from inhibiting

- Abstract
- ▼ Introduction
- ▼ Methods
- ▼ Results
- ▼ Discussion
- ▼ References

This Article

- [Abstract FREE](#)
- [Full Text \(PDF\)](#)
- [Alert me when this article is cited](#)
- [Alert me if a correction is posted](#)
- [Citation Map](#)

Services

- [Email this article to a friend](#)
- [Similar articles in this journal](#)
- [Similar articles in PubMed](#)
- [Alert me to new issues of the journal](#)
- [Download to citation manager](#)
- [Cited by other online articles](#)

Google Scholar

- [Articles by Zhang, C.](#)
- [Articles by Schmidt, J. T.](#)
- [Articles citing this Article](#)

PubMed

- [PubMed Citation](#)
- [Articles by Zhang, C.](#)
- [Articles by Schmidt, J. T.](#)



Content Types Subjects

English



Athens Authentication Point

Welcome!

To use the personalized features of this site, please **log in** or **register**.

If you have forgotten your username or password, we can **help**.

My SpringerLink

Marked Items

Alerts

Order History

Saved Items

All

Favorites

**Journal Article**

Effects of several 5'-carboxamide derivatives of adenosine on adenosine receptors of human platelets and rat fat cells

Journal Naunyn-Schmiedeberg's Archives of Pharmacology
 Publisher Springer Berlin / Heidelberg
 ISSN 0028-1298 (Print)
 1432-1912 (Online)
 Subject Biomedical and Life Sciences and Medicine
 Issue Volume 327, Number 1 / March, 1984
 DOI 10.1007/BF00504989
 Pages 36-42
 SpringerLink Sunday, December 12, 2004

Add to marked items

Add to shopping cart

Add to saved items

Recommend this article

D. Ukena¹, E. Böhme¹ and U. Schwabe¹

(1) Pharmakologisches Institut der Universität Heidelberg, Im Neuenheimer Feld 366, D-6900 Heidelberg 1, Federal Republic of Germany

Received: 13 February 1984 **Accepted:** 12 April 1984

Summary The effects of several 5'-carboxamide derivatives of adenosine on stimulatory (R_a) adenosine receptors of human platelets and inhibitory (R_i) adenosine receptors of rat fat cells have been compared. 5'-N-Cyclopropylcarboxamidoadenosine (CPCA) and 5'-N-ethylcarboxamidoadenosine (NECA) most potently inhibited ADP-induced aggregation of human platelets as shown by IC_{50} -values of 0.24 and 0.34

Find more options

 ...

- ☒ Within this issue
- ☐ Within this journal
- ☐ Within all content

Export this article

Export this article as RIS | Text

Text**PDF**

The size of this document is 820 kilobytes. Although it may be a lengthier download, this is the most authoritative online format.

Open: Entire document

Referenced by

2 newer articles

1. Cristalli, Gloria (2006) Highlights on the

$\mu\text{mol/l}$. 5'-N-Methylcarboxamidoadenosine (MECA; IC_{50} 0.81 $\mu\text{mol/l}$) and 5'-N-carboxamidoadenosine (NCA; IC_{50} 2.1 $\mu\text{mol/l}$) were less potent, whereas adenosine, 2-chloroadenosine and (-) N^6 -phenylisopropyladenosine [(-)PIA] exhibit IC_{50} -values of about 1.5 $\mu\text{mol/l}$. Nearly the same rank order of potency was obtained for stimulation of adenylate cyclase activity of platelet membranes and for inhibition of [^3H] NECA binding to human platelets. In order to examine the effects of the carboxamide analogues on R_i adenosine receptors of rat fat cells inhibition of lipolysis and adenylate cyclase were studied. (-)PIA was the most potent inhibitor of lipolysis as shown by an IC_{50} of 0.5 nmol/l, followed by CPCA (IC_{50} 1.1 nmol/l) and NECA (IC_{50} 1.3 nmol/l), whereas MECA (IC_{50} 17.9 nmol/l) and NCA (IC_{50} 20.1 nmol/l) were much less potent than NECA in inhibiting lipolysis. Similar results were obtained for inhibition of adenylate cyclase activity of fat cell membranes and for competition with [^3H]PIA binding to fat cell membranes. The relative potencies of the adenosine analogues at both receptor subclasses were calculated from the ratio of the IC_{50} -values for inhibition of platelet aggregation and of lipolysis. (-)PIA showed the highest selectivity for R_i receptors as indicated by a 2,900-fold lower IC_{50} for the antilipolytic than for the antiaggregatory effect. The R_a/R_i activity ratio for NECA was about 260, for CPCA 220, for NCA 105 and for MECA 45. These results indicate that all 5'-carboxamide adenosine derivatives are more potent agonists at R_i receptors than at R_a receptors. Since MECA has a higher

Development of A2A Adenosine Receptor Agonists and Antagonists. *ChemMedChem* [CrossRef]

- Gessi, S (2000) A2A adenosine receptors in human peripheral blood cells. *British Journal of Pharmacology* 129(1) [CrossRef]

[Ads by Google](#)

MECA Electronics, Inc.
RF/Microwave Coaxial Components N SMA BNC 7/16 DIN TNC, DC-18 GHz
www.e-meca.com

Adenosine Syringe Pump
Advanced Infusion 12S Pro. Perfect for pharm stress.
cnmtstore.com

Adenosine
Get the information you need to make an informed choice.
www.dlife.com

Fix Meca.exe
Download a Free Scan Tool & Fix Meca.exe Instantly!
www.TuneupAdvisor.com

Protein Characterization
Protein Solution's DynaPro Now available only through Wyatt!
www.wyatt.com

[Advertise on this site](#)

J Neurophysiol 78: 1965-1972, 1997;
0022-3077/97 \$5.00

The Journal of Neurophysiology Vol. 78 No. 4 October 1997, pp. 1965-1972
Copyright ©1997 by the American Physiological Society

NMDA-Independent LTP by Adenosine A₂ Receptor-Mediated Postsynaptic AMPA Potentiation in Hippocampus

Kofi Kessey¹ and David J. Mogul^{1, 2}

¹ Department of Neurobiology and Physiology and ² Department of
Biomedical Engineering, Northwestern University, Evanston, Illinois
60208

► ABSTRACT

Kessey, Kofi and David J. Mogul. NMDA-independent LTP by adenosine A₂ receptor-mediated postsynaptic AMPA potentiation in hippocampus. *J. Neurophysiol.* 78: 1965-1972, 1997. The role of adenosine A₂ receptors in normal synaptic transmission

and tetanus-induced long-term potentiation (LTP) was tested by stimulation of the Schaffer collateral pathway and recording of the field excitatory postsynaptic potential (EPSP) in the CA1 region of rat transverse hippocampal slices. Activation of adenosine A₂ receptors with the A₂ agonist N⁶-[2-(3,5-dimethoxyphenyl)-2-(2-methylphenyl)-ethyl]adenosine (DPMA; 20 nM) enhanced synaptic transmission during low-frequency test pulses (0.033 Hz). Paired stimulation before and during DPMA exposure indicated no paired-pulse facilitation as a result of A₂ activation, suggesting that enhancement was not a result of presynaptic modulation. DPMA enhanced the early phase α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) component of the EPSP. In contrast, DPMA had no effect on the N-methyl-D-aspartate (NMDA) component isolated using low extracellular Mg²⁺ and the AMPA receptor blocker 6-cyano-7-nitroquinoxaline-2,3-dione (20 μ M), indicating that the effects of A₂ activation on synaptic transmission were mediated by a postsynaptic enhancement of the AMPA response. Activation of adenosine A₂ receptors during a brief tetanus (100 Hz, 1 s) increased the level of LTP by 36% over that seen in response to a tetanus under control conditions. DPMA exposure after prior induction of LTP showed no additional potentiation, indicating that the mechanisms that contribute to both types of increases in synaptic transmission share a common mechanism. A slow onset NMDA-independent LTP could be induced by application of a tetanus during perfusion of DPMA with the NMDA blocker AP5 (50 μ M).

- Abstract
- ▼ Introduction
- ▼ Methods
- ▼ Results
- ▼ Discussion
- ▼ References

This Article

- [Abstract](#) FREE
- [Full Text \(PDF\)](#)
- [Alert me when this article is cited](#)
- [Alert me if a correction is posted](#)
- [Citation Map](#)

Services

- [Email this article to a friend](#)
- [Similar articles in this journal](#)
- [Similar articles in PubMed](#)
- [Alert me to new issues of the journal](#)
- [Download to citation manager](#)
- [Cited by other online articles](#)

Google Scholar

- [Articles by Kessey, K.](#)
- [Articles by Mogul, D. J.](#)
- [Articles citing this Article](#)

PubMed

- [PubMed Citation](#)
- [Articles by Kessey, K.](#)
- [Articles by Mogul, D. J.](#)